

Comparison of Supernovae datasets Constraints on Dark Energy

Zhang Cheng-wu, Xu Li-xin, Chang Bao-rong, and Liu Hong-ya
*School of Physics and Optoelectronic Technology,
 Dalian University of Technology, Dalian, 116024, P.R. China*

Cosmological measurements suggest that our universe contains a dark energy component. In order to study the dark energy evolution, we constrain a parameterized dark energy equation of state $w(z) = w_0 + w_1 \frac{z}{1+z}$ using the recent observational datasets: 157 Gold type Ia supernovae and the newly released 182 Gold type Ia supernovae by maximum likelihood method. It is found that the best fit $w(z)$ crosses -1 in the past and the present best fit value of $w(0) < -1$ obtained from 157 Gold type Ia supernovae. The crossing of -1 is not realized and $w_0 = -1$ is not ruled out in 1σ confidence level for the 182 Gold type Ia supernovae. We also find that the range of parameter w_0 is wide even in 1σ confidence level and the best fit $w(z)$ is sensitive to the prior of Ω_m .

PACS numbers: 98.80.Es, 98.80.-k, 95.36.+x

The universe is currently accelerating revealed by recent observations of type Ia supernovae (SNe Ia)[1, 2] and the acceleration has been attributed to a mysterious component dubbed dark energy. Although a cosmological constant Λ is the simplest candidate for dark energy and appears to explain current observations satisfactorily[3, 4], it suffers from fine tuning and coincidence problems[5]. These difficulties have lead to a large variety of proposed models with time-depended dark energy, such as quintessence[6], phantom[7], quintom[8], K-essence[9], tachyonic matter[10] and so on. In all the above models, the evolution history of the universe and the nature of dark energy strongly depend on the models, such as the dynamics of the extra energy component and the gravity theory used to solve the cosmological problem. However, there is another approach to study dark energy properties directly by observations in a model independent manner. In this method, the equation of state of dark energy $w(z) = p(z)/\rho(z)$ could be tested by cosmological observations to understand the gravitational properties of dark energy and its evolution. The aim of this paper is to compare the 157 Gold SNe Ia (SN157)[11] with the latest 182 Gold SNe Ia (SN182)[12] in constraining dark energy with a parameterized equation of state[13, 14, 15, 16]

$$w(z) = w_0 + w_1 \frac{z}{1+z} \quad (1)$$

where w_0 and w_1 are constant, z is redshift. Since the spatially flat universe $\Omega_m + \Omega_{DE} = 1$ agrees well with observations[3, 4], in this paper, we just consider the flat Friedmann-Robertson-Walker cosmology.

In a flat universe with Eq.(1), the Friedmann equation can be expressed as

$$H^2(z) = H_0^2 [\Omega_m (1+z)^3 + \Omega_{DE} (1+z)^{3(1+w_0+w_1)} e^{\frac{-3w_1 z}{(1+z)}}] \quad (2)$$

Then the knowledge of Ω_m and $H(z)$ is sufficient to determine $w(z)$.

Supernovae observations provide the apparent magnitude $m(z)$ at peak brightness after implementing correction for galactic extinction, K-correction and light curve

width-luminosity correction. $m(z)$ is related to the luminosity distance $d_L(z)$ through

$$\begin{aligned} m_{th}(z) &= 5 \log_{10}(D_L(z)) + M + 5 \log_{10}\left(\frac{H_0^{-1}}{Mpc}\right) + 25 \\ &= 5 \log_{10}(D_L(z)) + M + 42.38 - 5 \log_{10} h \end{aligned} \quad (3)$$

where $H_0 = h \text{ km} \cdot \text{s}^{-1} \cdot \text{Mpc}^{-1}$, parameter M is the absolute magnitude which is assumed to be constant and $D_L(z)$ is defined as

$$\begin{aligned} D_L(z) &= H_0 d_L(z)/c \\ &= (1+z) \int_0^z \frac{H_0 dz'}{cH(z')} \end{aligned} \quad (4)$$

Supernovae data points are given in terms of the distance modulus

$$\mu_{obs}(z_i) \equiv m_{obs}(z_i) - M \quad (5)$$

and theoretical model parameters are determined by minimizing the quantity

$$\chi^2 = \sum_{i=1}^N \frac{(\mu_{obs}(z_i) - \mu_{th}(z_i))^2}{\sigma_{(obs;i)}^2} \quad (6)$$

where $\sigma_{(obs;i)}^2$ are the errors due to flux uncertainties, intrinsic dispersion of SNe Ia absolute magnitude and peculiar velocity dispersion respectively. These errors are assumed to be gaussian and uncorrelated. The theoretical distance modulus is defined as

$$\begin{aligned} \mu_{th}(z_i) &\equiv m_{th}(z_i) - M \\ &= 5 \log_{10}(D_L(z)) + 42.38 - 5 \log_{10} h \end{aligned} \quad (7)$$

and μ_{obs} is given by supernovae dataset.

Our calculations to limit parameters are based on the Markov Chain Monte Carlo program CosmoMC[17]. We modified the code to make equation of state w be redshift-depended instead of constant -1 and add parameter w_0 and w_1 as slow parameters to the cosmological parameters space. Using the maximum likelihood method with




FIG. 1: The contours show 2-D marginalized 1σ and 2σ confidence limits in the (w_1, w_0) , (w_1, Ω_m) , (Ω_{DE}, Ω_m) , (w_0, Ω_m) plane from SN157.




FIG. 2: The contours show 2-D marginalized 1σ and 2σ confidence limits in the (w_1, w_0) , (w_1, Ω_m) , (Ω_{DE}, Ω_m) , (w_0, Ω_m) plane from SN182.




FIG. 3: The best fits of $w(z)$ with 1σ errors (shaded region). Ω_m is set to the best values obtained from SN157 or SN182.




FIG. 4: The best fits of $w(z)$ with 1σ errors (shaded region) with a prior $\Omega_m = 0.27$.

TABLE 1. The minimum χ^2 and mean fit values from SN157 and SN182.

data	χ^2_{min}	w_0	w_1	Ω_m
SN157	89.947	-1.856 ± 0.519	-0.015 ± 0.269	0.437 ± 0.073
SN182	81.523	-1.637 ± 0.496	-0.180 ± 0.081	0.435 ± 0.058

SN157 and SN182, we obtained confidence level of equation of state and matter density. The contour plots are show in Fig.1 and Fig.2, the mean fit values, 1σ constraints on parameters w_0 , w_1 and Ω_m with standard deviations are list in Table 1. In Fig.1 we show results for the parameterized equation of state for the SN157 dataset. We find the best fit value of (w_1, w_0, Ω_m) to be $(0.335, -2.373, 0.485)$ and see that the (w_1, w_0) panel has two island separately as well as the (w_1, Ω_m) panel which means that the SN157 dataset can not give a fine constraint on w_1 . Fig.2 shows the results calculated by the SN182 dataset. The best fit value of (w_1, w_0, Ω_m) is $(-0.244, -1.728, 0.462)$.

Fig.3 shows 1σ errors of the best fit $w(z)$ calculated using the covariance matrix[18]. The best fit $w(z)$ constrained by SN157 can cross -1 in the evolution and in 1σ confidence level, the LCDM parameter ($w_0 = -1, w_1 = 0$) do not coincide with the best fit value of $w(z)$ at present. But the best fit dynamical $w(z)$ obtained from SN182 does not cross -1 for the whole redshift range $0 < z < 1.75$, $w_0 = -1$ is in 1σ confidence contour and near the boundary. During our calculations, we also find that the best fit $w(z)$ and its errors are sensitive to the value of Ω_m . When we set $\Omega_m = 0.27$ as a prior instead of the best fit value mentioned before, both of the best fit $w(z)$ can cross -1 and its 1σ error region shrink sharply as showed in Fig.4.

In summary, we have provided a comparative analysis on constraining parameters w_1, w_0 , Ω_m and the properties of dark energy with two cosmological observations: SN157 and SN182. It shows that the SN157 dataset can not give a fine constraint on w_1 and the best fit $w(z)$ can cross -1 in the evolution, but the SN182 dataset gives contrary result. We also find that the range of parameter w_0 is wide even in 1σ confidence level and the best fit $w(z)$ is sensitive to the prior of Ω_m . In the future, we can combine other cosmological observations, such as Cosmic Microwave Background anisotropies[4], the SDSS baryon acoustic peak[19], the cluster baryon fraction (CBF)[20] or the linear perturbations growth rate at $z = 0.15$ obtained from the 2dF galaxy redshift survey (PGR)[21] to constrain the matter density and the evolution of dark energy. We hope that the best fit values of cosmological parameters and the confidence level could be enhanced which will help us to understand the nature of dark energy.

Acknowledgements We acknowledge the using of CAMB[22] and CosmoMC[23] for the numerical calculations. Chengwu Zhang thanks the helpful discussion with

Prof. Antony Lewis[24] and suggestions from Dr. Haitao Cui. This work was supported by NSF (10573003), NBRP (2003CB716300) of P.R. China. The research of

Lixin Xu was also supported in part by NSF (10647110) and DUT 893321.

-
- [1] Riess A G et al 1998 *Astron. J.* **116** 1009
astro-ph/9805201
 - [2] Perlmutter S et al 1999 *Astrophys. J.* **517** 565
astro-ph/9812133
 - [3] Spergel D N et al 2003 *Astrophys. J. Suppl.* **148** 175
astro-ph/0302209
 - [4] Spergel D N et al 2006 *arXiv* astro-ph/0603449
 - [5] Peebles P J E and Ratra B 2003 *Rev. Mod. Phys.* **75** 559
astro-ph/0207347
 - [6] Zlatev I, Wang L M and Steinhardt P J 1999 *Phys. Rev. Lett.* **82** 896
 - [7] Caldwell R R, Kamionkowski M and Weinberg N N 2003 *Phys. Rev. Lett.* **91** 071301
 - [8] Feng B, Wang X L and Zhang X M 2005 *Phys. Lett. B* **607** 35
 - [9] Armendariz-Picon C, Damour T and Mukhanov V 1999 *Phys. Lett. B* **458** 209
 - [10] Padmanabhan T 2002 *Phys. Rev. D* **66** 021301
 - [11] Riess A G et al 2004 *Astrophys. J.* **607** 665
astro-ph/0402512
 - [12] Riess A G et al 2006 *arXiv* astro-ph/0611572
 - [13] Chevallier M and Polarski D 2001 *Int. J. Mod. Phys. D* **10** 213 gr-qc/0009008
 - [14] Linder E V 2003 *Phys. Rev. Lett.* **90** 091301
astro-ph/0208512
 - [15] Choudhury T R and Padmanabhan T 2005 *Astron. Astrophys.* **429** 807 astro-ph/0311622
 - [16] Nesseris S and Perivolaropoulos L 2005 *Phys. Rev. D* **72** 123519 astro-ph/0511040
 - [17] Lewis A and Bridle S 2002 *Phys. Rev. D* **66** 103511
 - [18] Alam U et al 2004 *ArXiv* astro-ph/0406672
 - [19] Eisenstein D J et al 2005 *Astrophys. J.* **633** 560
 - [20] Allen S W et al 2004 *Mon. Not. R. Astron. Soc.* **353** 457
astro-ph/0405340
 - [21] Verde L et al 2002 *Mon. Not. R. Astron. Soc.* **335** 432
astro-ph/0112161
 - [22] [Http://camb.info](http://camb.info)
 - [23] [Http://cosmologist.info/cosmomc](http://cosmologist.info/cosmomc)
 - [24] [Http://cosmocoffee.info](http://cosmocoffee.info)